



ISSN 2456-3110

Vol 7 · Issue 8

September 2022

Journal of  
**Ayurveda and Integrated  
Medical Sciences**

*www.jaims.in*

JAIMS

An International Journal for Researches in Ayurveda and Allied Sciences



**Maharshi Charaka**  
Ayurveda

Indexed

# A critical study on Burn Injury (*Dagdha Vrana*) with special reference to *Sushruta Samhita*

Anagha TS<sup>1</sup>, KM Sweta<sup>2</sup>, Dhyan Surendranath<sup>3</sup>

<sup>1</sup>Post Graduate Scholar, Department of Shalya Tantra, Sri Sri College of Ayurvedic Science and Research Hospital, Bengaluru, Karnataka, India.

<sup>2</sup>Professor & HOD, Department of Shalya Tantra, Sri Sri College of Ayurvedic Science and Research Hospital, Bengaluru, Karnataka, India.

<sup>3</sup>Associate Professor, Department of Shalya Tantra, Sri Sri College of Ayurvedic Science and Research Hospital, Bengaluru, Karnataka, India.

## ABSTRACT

Burns are global public health problem, accounting for an estimated death 180000 annually and the estimated annual burn incidence in India is approximately 6 -7 million. In Ayurveda, burn injury has been dealt in the name of *Dagdha Vrana*. The concept of *Dagdha Vrana* was found in different treatises, In *Sushruta Samhita* details regarding classification, clinical features and detailed management according to types are described. In Ayurveda, therapeutic burns also have been described. This article compiles with classification, clinical features and treatment of burns injury and *Dagdha Vrana* with special reference to *Sushruta Samhita*.

**Key words:** *Dagdha Vrana*, Burn injury, *Sushruta Samhita*.

## INTRODUCTION

Burn injuries are an under-appreciated trauma that can affect anyone, anytime and anywhere.<sup>[1]</sup> A burn is a wound in which there is coagulative necrosis of the tissue. Also defined as damage to the skin or deeper tissues caused by sun, hot liquids, fire, electricity, radiations or chemicals is a burn.<sup>[2]</sup> Ayurveda which is an ancient science of treatment where the description of types, clinical features, treatments and complications of *Dagdha Vrana* having similarity to

burn injury described in modern medicine. *Dagdha Vrana* (burn wound) which can be accidental or may be caused during para surgical procedures like *Agnikarma*. *Acharya Sushruta*, described *Dagdha Vrana* in the chapter '*Agnikarma Vidhi Adhyaya*' in *Sutrasthana*. In this *Adhyaya*, *Acharya* explains the clinical features, types and treatment protocol for individual type of burns.<sup>[3]</sup>

### Types of Burn Injury

#### a) Thermal Injury<sup>[4]</sup>

Thermal injury are mainly scalds, flame, flash burns

**Scalds:** It is mainly due to spillage of hot liquids having temperature > 65°C. It is capable of creating a deep partial-thickness or full-thickness burn in 3 seconds of contact. The same burn occurs in 1 second if the temp is around 70°C. In children under 8 years of age, the most common burns are scalds, usually from the spilling of hot liquids.

**Flame Burns:** It is the second most common mechanism of thermal injury. Damage causing mainly from superheated oxidized air.

### Address for correspondence:

Dr. Anagha TS

Post Graduate Scholar, Department of Shalya Tantra, Sri Sri College of Ayurvedic Science and Research Hospital, Bengaluru, Karnataka, India.

E-mail: anagha123suresh@gmail.com

Submission Date: 15/07/2022 Accepted Date: 21/08/2022

### Access this article online

#### Quick Response Code



Website: [www.jaims.in](http://www.jaims.in)

Published by Maharshi Charaka  
Ayurveda Organization, Vijayapur,  
Karnataka (Regd) under the license  
CC-by-NC-SA

E.g.: Includes improper use of flammable liquids, motor vehicle collisions, and ignition of clothing by stoves. In older children and adults, the most common burns are flame-related, usually the result of house fires, and the ill-advised use of flammable liquids.

**Flash Burns:** These burns are typically epidermal or partial thickness and are caused mainly by Explosions of natural gas, propane, butane, petroleum distillates, alcohols, and other combustible liquids

#### b) Electrical Injury<sup>[5]</sup>

Injury severity is determined by the voltage of the source and the resistance of the victim. Tissue damage in electrical injury occurs when electrical energy is converted to thermal energy. The resulting injury is a thermal burn. Electrical burns are always deep burns, and wound present at point and entry and exit. The peculiarity of electrical burns is that it causes minimal destruction of the skin. The skin is involved at 2 points - at the point of contact with the electrical source and at the site of exit at which the patient is grounded. Depending on the voltage it is again divided into,

**Low Voltage Current:** Where there is direct injury at point of contact and there will be local coagulation and necrosis.

**High-Voltage Current:** Where there is minimum destruction of skin but damages tissues that conduct the electricity through the body. Tissue damage in electrical injury occurs when electrical energy is converted to thermal energy. The resulting injury is a thermal burn.

#### c) Chemical Burn<sup>[6]</sup>

Chemical burn is caused by contact with strong acid / base.

Acids "tan" the skin, creating an impermeable barrier of coagulation necrosis and limits further penetration. E.g.: Sulphuric acid, Nitric acid

Alkalis combine with cutaneous lipids, and continue dissolving the skin until they are neutralized. Eg: Ammonia, Cement

Chemical burns are typically associated with industrial accidents, assaults, or the improper use of harsh

household solvents and cleaners. Chemical burns because progressive damage and injury until the chemicals are inactivated by reaction with tissues or diluted by therapeutic irrigation. The depth of the burn can only be assessed after removal of slough.

#### d) Radiation Burn<sup>[7]</sup>

Radiation burns are generally a complication of therapeutic administration and are result of free oxygen radical formation. Radiations are mainly classified as nonionizing radiation (Radio wave, Microwave, IR, UV) and ionizing radiation (x ray, Gamma rays). Because of radiation there are superficial and deeper injuries. In radiation burn two types of radiodermatitis are seen that is acute radio dermatitis and chronic radiodermatitis.

**Acute Radiodermatitis** which presents with acute inflammation with erythema, varying degrees of oedema and exfoliation.

**Chronic Radiodermatitis** may occur if small doses of irradiation are given for too long time, in this condition the skin shows irregular pigmentation or depigmentation. And Long span of exposure will lead to indolent ulcers.

#### e) Cold Burn<sup>[8]</sup>

Cold burns are caused by exposure to cold which include freezing injury [frost bite], non-freezing cold injury [chill blain, trench foot]. Clinical features of frostbite are described by various degrees.

First Degree Frostbite: there will be hyperemia and oedema of the skin.

Second Degree Frostbite causes hyperemia, vesicle formation and partial thickness necrosis of the skin.

Third degree frostbite: causes necrosis of the entire skin thickness and may extend to subcutaneous tissue.

Fourth degree frostbite: causes necrosis of full thickness of the skin including subcutaneous tissue, muscle and bone. This leads to gangrene of the affected part.

Trench Foot: Due to prolonged exposure to cold weather with circulatory disturbances.

Chill Blain: Due to cold weather there will be localized erythema in extremities.

According to Sushruta Acharya<sup>[9]</sup>

तत्र, स्निग्धं रूक्षं वाऽ(चा)श्रित्य द्रव्यमग्निर्दहति; अग्नि सन्तप्तो हि स्नेहः सूक्ष्मसिरानुसारित्व त्वगादीन् अनुप्रविश्याशु दहति; तस्मात् स्नेहदग्धेऽधिका रुजो भवन्ति । (Su. Su. 12/15)

Sushruta Acharya classified as Snigdha and Ruksha.

Heat which is produced by either Ruksha or Snigdha substances causes Dagdha. Burn produced due to Snigdha Dravyas are more painful, as the Sneha Dravya has the property of penetrating the minute pores and causes deeper and more painful burns.

### Classification of Burns

#### 1. Depending on thickness of Skin Involved<sup>[8]</sup>

Depending on thickness of skin involved, they are classified into 4.

**First Degree:** In which Epidermis looks red and it is painful, absence of blisters. It heals rapidly in 5-7days without scarring and shows capillary filling.

**Second Degree:** the affected area is mottled, red, and painful, with blisters Heals by 14-21 days. It is again classified into Superficial 2nd degree burns which heal by causing pigmentation and Deep 2<sup>nd</sup> degree burns which cause scarring. Sensation will be present.

**Third Degree:** The affected area is charred, parchment like painless and insensitive. There will be eschar formation and re-epithelization occurs from wound edge.

**Fourth Degree:** It involves the underlying tissues including muscles, bones.

#### 2. Depending on Severity of Burn<sup>[7]</sup>

Depending on severity of burns, it is classified as

**Superficial partial-thickness** burns which include superficial 2<sup>nd</sup> degree burns, involving the upper layers of dermis, form blisters with fluid collection. When blisters are removed, the wound is pink, wet; hypersensitive, and burns blanch with pressure.

**Deep partial-thickness burns** extend into the reticular layers of the dermis. There will be blister formation, but the wound surface is usually a mottled pink-and-white color immediately after the injury because of the varying blood supply to the dermis

**Full-thickness burns** including 3<sup>rd</sup> and 4<sup>th</sup> degree burns involve all layers of the dermis and can heal only by wound contracture, and epithelialization occurs from the wound margin.

#### 3. Depending on the percentage of Burns<sup>[7]</sup>

Percentage of burns are mainly classified into mild, moderate, severe.

**Mild:** Includes Partial thickness burns which is < 15% of total body surface in adults or <10% in children. When it is Full thickness which include < 2% of body surface area.

**Moderate:** Includes Second degree of burns which involved 15 -20% in adults (10-20% in children) and 3<sup>rd</sup> degree burns between 2-10%. Not involving eyes, hands, ears, face and perineum.

**Severe:** Includes 2<sup>nd</sup> degree burns which is >25% in adults and >20% in children. Also 3<sup>rd</sup> degree burns of 10% or more. Involving eyes, ears, feet and hand.

According to Sushrut Acharya<sup>[9]</sup>

तत्र प्लुष्टं दुर्दग्धं सम्यग्दग्धमतिदग्धं चेति चतुर्विधमग्निदग्धम् । (Su.Su.12/16)

The types according to Acharya Sushruta,

- प्लुष्टं
- दुर्दग्धं
- सम्यक् दग्धम्
- अतिदग्धम्

**Plusta Dagdha<sup>[9]</sup>**

तत्र यद्विवर्णं प्लुष्यतेऽतिमात्रं तत् प्लुष्टं (su.su,12/16)

There will be discoloration of the skin of affected part along with burning sensation. Dalhanacharya specifies that Plusta Dagdha is the one causing discoloration and Daha but no Spotha. It can be co-related to 1<sup>st</sup> degree burns.

**Durdagdha<sup>[9]</sup>**

यत्रोत्तिष्ठन्ति स्फोटास्तीव्राश्लोषदाहरागपाकवेदनाश्चिराच्चोप  
शाम्यन्ति तद्दुर्गंधं (Su.Su,12/16)

Burns presenting with blisters, and different types of pain, burning, redness, inflammation. Also takes a long time to subside. Can be correlated to 2<sup>nd</sup> degree burns.

**Samyak Dagdha<sup>[9]</sup>**


सम्यग्दग्धमनवगाढं तालवर्णं सुसंस्थितं पूलक्षणयुक्तं च  
(Su.Su.12/16)

Burns which are not deeply seated nor superficial. Having the color of *Tala Phala* - Asian palm fruit. Tissues will be in their own site. And having signs and symptoms described earlier.

**Atidagdha<sup>[9]</sup>**

अतिदग्धे मांसावलम्बनं गात्रविक्षेपः सिरास्नायुसन्ध्यस्थि  
व्यापादनमतिमात्रं ज्वरदाहपिपासामूर्च्छाश्लोषद्रवा भवन्ति,  
व्रणश्चास्य चिरेण रोहति, रूढश्च विवर्णो भवति ||  
(Su.Su.12/16)

The injury affecting the whole tissue as skin subcutaneous tissue, muscles, vessels, tendons, ligaments and exposing to other deeper structure. The patients having high grade fever, burning pain, increase thirst, features of dehydration. The wound healing delayed with discoloration of affected part. The condition is almost similar to advanced 3<sup>rd</sup> degree and 4<sup>th</sup> degree burn.

**Assessment of Burn<sup>[10,11]</sup>****1. Wallace rule of 9**

In this method, every part of body is assessed. For Children which have a relatively larger portion of the body surface area in the head and neck, which is compensated for by a relatively smaller surface area in the lower extremities. Infants have 21% of the TBSA in the head and neck and 13% in each leg, which incrementally approaches the adult proportions with increasing age.

- Lund and Browder chart:** Each part is individually assessed for burns. More useful in children taking in consideration the age and BSA of the patient.
- Palmar method:** Patient own hand is used to know the extent of burn. The entire palm of person who is burned is considered as 0.8 % in adults and 1 % in children, and using the palm to measure the body surface.
- Sage diagram:** Software program which helps in calculating the extent of burn and also the fluid resuscitation levels

**Pathophysiology<sup>[12]</sup>**

Due to injury, Heat causes coagulation necrosis of skin and subcutaneous tissue which lead to release of vasoactive peptides. Because of this there will be altered capillary permeability which further leads to Loss of fluid and Severe Hypovolemia. Hypovolemia leads to Decreased cardiac output which leads to altered pulmonary resistance, Decreased Myocardial function and Decreased Renal blood flow (Oliguria) which further leads to SIRS and MODS.

**Management of Burns<sup>[12,13]</sup>****Initial Management**

- Clothing should be removed initially.
- Cooling of the part by running water for 20 mins (should not use cold water, to prevent from Hypothermia)
- Chemoprophylaxis - tetanus toxoid (0.5ml), antibiotics, local antiseptics
- Comforting with Sedation and Analgesics.

**Fluid Resuscitation**

First 24 hours, Crystalloids should be used because it is easily permeable through capillaries and replace blood volume. After 24 hours up to 48 hours colloids should be used to compensate plasma loss (they are retained in intravascular compartment)

Ryle's tube insertion: Should be done initially for aspiration then feeding (burns <15%)

Total parenteral nutrition (TPN) using CVP line, lipids and vitamins for faster recovery

Urinary catheterization - to maintain fluid volume - 30-40ml/hr

Vitals : hourly pulse, SPO2, BP, electrolyte analysis, PO2, PCO2, blood urea etc.

Frequent total count and platelet count - identify sepsis along with fever, tachycardia etc.

**Local Management<sup>[14]</sup>**

Regular dressing should be done with Paraffin gauze, Hydrocolloids etc. under suitable anesthesia.

**Open Method:** Application of silver sulfadiazine without any dressings, used commonly in burns of face, head and neck.

**Closed Method:** Dressings done to soothen and to protect the wound, to reduce the pain

**Tangential Excision:** Excision of burn wound with skin grafting can be done within 48 hours in patients with less than 25% burns. Done in deep dermal burn wherein dead dermis is removed layer by layer until fresh bleeding occurs. Later skin grafting is done. It reduces the chance of secondary infection, the hospital stays, and formation of hypertrophic scar or contracture.

**Treatment according to Sushruta Acharya****Plusta Dagda Chikitsa<sup>[15]</sup>**

प्लुष्टस्याग्निप्रतपनं कार्यमुष्णं तथौषधम् ।  
शरीरे स्विन्नभूयिष्ठे स्विन्नं भवति शोणितम् ॥  
प्रकृत्या ह्युदकं शीतं स्कन्दयत्यतिशोणितम् ।  
तस्मात् सुखयति ह्युष्णं ननु शीतं कथञ्चन ॥  
(Su.Su.12/19-21)

Acharya Sushruta has mentioned the use of warm medication for the treatment of *Plusta Dagda*, so as to prevent *Skandana* of the *Sonita*.

**Durdagda Chikitsa<sup>[16]</sup>**

शीतामुष्णां च दुर्दग्धे क्रियां कुर्याद्विषक् पुन ।

घृतालेपनसेकांस्तु शीतानेवास्य कारयेत् । (Su.Su.12/22)

Acharya Sushruta mentions the use of both warm and cold medication in this type of burn. Acharya Dalhana, commented on it saying that; if it is a deep burn then internal medicine having cold active principles should be used and if it is a superficial burn then drugs having warm active principles to be used. For external application - *Ghrita Lepa*, *Seka* and cold application is used.

**Samyak Dagda Chikitsa<sup>[16]</sup>**

सम्यग्दग्धे तुगाक्षीरीप्लक्षचन्दनगैरिकैः।

सामृतैः सर्पिषा स्निग्धैरालेपं कारयेद्विषक् ॥

ग्राम्यान्पौदकैश्चैनं पिष्टैर्मासैः प्रलेपयेत् ।

पित्तविद्राधिवच्यैर्न सन्ततोष्माणमाचरेत् ॥ (Su.Su.12/23-24)

As per Acharya Sushruta advises *Lepa* prepared of *Tugakshiri*, *Plaksha*, *Chandan*, *Amrita* & *Gairik* with *Gritha* can be used as local application. Paste of flesh of animals like horse, pig etc. may also be used as *Lepa*. If there is continuous burning sensation then line of treatment should be same as *Pittaj Vidradhi*.

**Atidagda Chikitsa<sup>[16]</sup>**

अतिदग्धे विशीर्णानि मांसान्युद्धृत्य शीतलाम् ।

क्रियां कुर्याद्विषक् पश्चाच्छालितण्डुलकण्डनैः ॥

तिन्दुकीत्वक्कपालैर्वा घृतमिश्रैः प्रलेपयेत् ।

व्रणं गुडूचीपत्रैर्वा छादयेदथवौदकैः ॥

क्रियां च निखिलां कुर्याद्विषक् पित्तविसर्पवत् । (Su.Su.12)

As per Acharya Sushruta, *Atidagda* is similar to advanced third / fourth degree burn, management included surgical removal of dead tissue followed by medical treatment by above mentioned drugs as local application. Acharya Sushruta indicated to avoid

bandages on affected part, the area should be covered with wet *Guduchi*, *Kamal Patra* etc. the rest of treatment should be same as *Pittaj Visarpa* (erysipelas). *Gauryadi Ghrita* for local application.

*Acharya Sushruta* has specially mentioned regarding Burns that caused by the *Vaidya* while performing *Agnikarma* for therapeutic reasons. They are;

#### *Twak Dagda*<sup>[12]</sup>

शब्दप्रादुर्भावो दुर्गन्धता त्वक्सङ्कोचश्च ॥

If the burn caused in *twak* the symptoms includes Charring sound, Foul smell and Shrinking of skin.

#### *Mamsa Dagda*<sup>[12]</sup>

कपोतवर्णताऽल्पश्वयथुवेदना शुष्कसङ्कुचितव्रणता च मांस दग्धे ॥

In *Mamsa Dagda*, there will be less inflammation, Discoloration, Shrinking and contraction of muscles.

#### *Sira and Snayu Dagdha*<sup>[12]</sup>

कृष्णोन्नतव्रणता स्रावसन्निरोधश्च सिरास्नायुदग्धे ॥

In *Sira Snayu Dagda*, there will be blackish discoloration, Elevated swelling and Excessive exudation.

#### *Sandhi Dagdha*<sup>[12]</sup>

रूक्षारुणता कर्कशस्थिरव्रणता च सन्ध्यस्थिदग्धे ॥

*Sandhi Dagda Lakshanas* include Dryness, Redness, Rough texture and Formation of wound.

### DISCUSSION

*Dagdhavrana* is a concept found in Ayurvedic treatise, most notably in *Sushruta Samhita*, the pioneering book on Indian surgery. *Acharya Sushruta* has mentioned the importance of understanding the intensity of burn based on the type of heat causing the burn. In *Agnikarma*, *Vaidykruta Dagdha* are mentioned as scales that indicate the depth of burn. Where in detailed explanation regarding the appearance and the symptoms of the burn in relation to the layers of the body is given.

*Sushruta Acharya* explains *Ushna Chikitsa* as the treatment principle for *Plushta Dagda* and *Seetopachara* is strictly contraindicated here, as it causes *Skandathwa* of *Shonita* which prevents dissipation of heat and traps heat locally causing further tissue damage. *Swedana* causes *Vilayana* of *Shonita* and improves local circulation thus dissipates heat prevents further tissue damage. *Acharya Sushruta* mentioned regarding *Bandhana* in *Vranalepanabandhana Vidhi Adhyaya*, in that the contraindications of *Bandhana*, included *Dagdha Vrana* also. This can be understood as *Badhana* done in an acute stage of burn injury it will in turn cause *Mamsapaka* and got worsen the condition. The current understanding of allopathic science also supports this.

### CONCLUSION

In regard to etiology, clinical features, degree of burn, types and treatment modalities described by *Acharya Sushruta* resemble those described in the modern system of medicine.

### REFERENCES

1. Makhija L, Bajaj S, Gupta J. National programme for prevention of burn injuries. *Ind. J. Plast. Surg.* 2010;43(3):6 [Internet], [cited 2020 Apr 10] Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3038407/>.
2. Sriram Bhat M. *SRB's Manual of Surgery. Burns: chapter 1[i].* 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 128.
3. *Susruta, Susruta Samhita, English translation by Srikantamurthy.* 1st ed. Suthrasthana, Ch-12. Varanasi: Chaukambha visvabharati; 2018. Vol-1.p.71
4. F.Charles et al, *Schwartz's principles of surgery, 8th Ed, Ch-8. McGraw hill's Access Medicine.* p.209
5. Sriram Bhat M. *SRB's Manual of Surgery. Burns: chapter 1[i].* 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 130.
6. F.Charles et al, *Schwartz's principles of surgery, 8th Ed, Ch-8. McGraw hill's Access Medicine.* p.212

7. Sriram Bhat M. SRB's Manual of Surgery. Burns: chapter 1[i]. 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 134.
8. Sriram Bhat M. SRB's Manual of Surgery. Burns: chapter 1[i]. 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 136.
9. Susruta, Susruta Samhita, English translation by Srikantamurthy. 1st ed. Suthrasthana, Ch-12. Varanasi: Chaukambha visvabharati; 2018. Vol-1.p.73
10. Sriram Bhat M. SRB's Manual of Surgery. Burns: chapter 1[i]. 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 137.
11. F.Charles et al, Schwartz's principles of surgery, 8th Ed,Ch-8.McGraw hill's Access Medicine.p.225
12. Das.S.A concise Textbook of Surgery,10<sup>th</sup> edition.Kolkata:Dr.S.Das.13;2018.P.61-79
13. Sriram Bhat M. SRB's Manual of Surgery. Burns: chapter 1[i]. 4th ed. New Delhi: Jaypee Brothers Medical Publishers; 2013. p. 139.
14. F.Charles et al, Schwartz's principles of surgery, 8th Ed,Ch-8.McGraw hill's Access Medicine.p.235
15. Susruta, Susruta Samhita, English translation by Srikantamurthy. 1st ed. Suthrasthana, Ch-12. Varanasi: Chaukambha visvabharati; 2018. Vol-1.p.74.
16. Susruta, Susruta Samhita, English translation by Srikantamurthy. 1st ed. Suthrasthana, Ch-12. Varanasi: Chaukambha visvabharati; 2018. Vol-1.p.75.

**How to cite this article:** Anagha TS, KM Sweta, Dhyan Surendranath. A critical study on Burn Injury (Dagda Vrana) with special reference to Sushruta Samhita. J Ayurveda Integr Med Sci 2022;8:86-92.

**Source of Support:** Nil, **Conflict of Interest:** None declared.

\*\*\*\*\*