



ISSN 2456-3110

Vol 2 · Issue 6

Nov-Dec 2017

Journal of
**Ayurveda and Integrated
Medical Sciences**

www.jaims.in

JAIMS



Charaka
Publications

Indexed

Anti-inflammatory effect of an Ayurveda polyherbal topical application and exploration of its mechanism of action

Sitaram Ahalya,¹ B. A. Venkatesh,² R. Vijayasarithi,³ Tirumalapura Vijayanna Shalini⁴

¹Joint Director, Dept. of AYUSH, Bangalore, ²Principal, Sri Sri College of Ayurvedic Medicine & Research Hospital, Bangalore, ³Retired Professor & H.O.D, Dept. of PG Studies in Shalya Tantra, Government Ayurveda Medical College, Bangalore, ⁴Research Associate, MS Ramaiah Indic Center for Ayurveda & Integrative Medicine, Bangalore, Karnataka, India.

ABSTRACT

Background: Topical application of herbs is one of the recommended treatment modalities for Osteoarthritis (OA) in Ayurveda (Indian system of medicine). The current study intended to evaluate the anti-inflammatory effect of a polyherbal powder used as *Upanaha* (poultice) for OA by in vitro and in vivo techniques. **Materials and Methods:** The polyherbal formulation *Upanaha Choornam* (UC) was sourced from *Vaidyaratnam Oushadhasala* (P) Ltd., Thrissur, India. Changes in the secretion of TNF- α and NO and expression of Cox-2 genes were evaluated by semi quantitative PCR activity to establish anti-inflammatory action in vitro. Macrophages and connective tissue of mice were used as media for the former two experiments and only macrophages for the latter. In vivo anti-inflammatory activity was evaluated by TPA induced ear oedema in Swiss Albino mice (n=24), divided into 4 groups as Group I - saline treatment, Group II - Indomethacin treatment, and Groups III and IV treated with 30% and 60% of UC respectively. **Results:** In the in vitro study, UC at 1000 μ g/ml and 500 μ g/ml upregulated the COX-2 level by 0.08 and 0.03 folds respectively as compared to control. Release of TNF- α , and NO in LPS-induced RAW cells were significantly inhibited in a dose dependent manner. The TPA induced ear oedema significantly reduced in Groups III and IV (F=1250, p<0.001) **Conclusion:** The current study demonstrates the safety and anti-inflammatory activity of a polyherbal formulation *Upanaha Choornam* as a topical application. This indicates the potential of select herbs in managing degenerative conditions like OA.

Key words: Osteoarthritis, Upanaha, Sandhigata Vata, Poultice, Anti-Inflammatory Activity.

INTRODUCTION

Ayurveda (Indian system of medicine) emphasises three major lines of treatment viz, internal medications and diet (*Antah Parimarjana*), external

therapies or topical applications (*Bahir Parimarjana*) and surgical/para-surgical procedures (*Shastra Pranidhana*) (C.S.Su11/55). These treatments are used in various combinations to provide optimal benefit in diverse disease conditions. Internal medications receive maximum attention for research studies, while the same cannot be said about external therapies. Osteoarthritis (OA), regarded as *Sandhigata Vata* in Ayurveda is a chronic degenerative condition where in topical application of herbs in the form of paste (*Kalka*), powder (*Churna*), oil (*Taila*) are widely prescribed.^[7] *Upanaha Churna* is one such polyherbal formulation, referenced from *Ashtanga Hridaya*, commonly prescribed for topical application in *Sandhigata Vata*.^[12] It consists of herbs *Acorus calamus* Linn. (*Vacha*), *Anethum sowa* Roxb.

Address for correspondence:

Sitaram Ahalya

Joint Director, Department of AYUSH, Dhanvantari road, Bangalore, Karnataka, India.

E-mail: ahalyasharma@yahoo.com

Submission Date : 12/11/2017 Accepted Date: 23/12/2017

Access this article online

Quick Response Code



Website: www.jaims.in

DOI: [10.21760/jaims.v2i06.10924](https://doi.org/10.21760/jaims.v2i06.10924)

(*Shatahwa*), *Cedrus deodara* Roxb. Ex.lamb (*Devadaru*), *Pluchea lanceolata* DC Heirn /Alpania galangal Lance. (*Rasna*), *Ricinus communis* Linn. (*Eranda*), *Nardostachys jatamansi* DC Heirn (*Jata*), Rock salt (*Saindhava Lavana*), grains like *Sesamum indica* Linn. (*tila*), *Brassica campestris* Linn. (*Sarshapa*), *Dolichus biflorus* Linn. (*Kulattha*), *Phaseolus mungo* Linn. (*Mudga*), *Phaseolus vulgaris* Linn. (*Masha*), *Oryza sativa* Linn. (*Shashtika*), *Panicum sumatrense* Roth ex Roem. & Schult. (*Shyamaka*), *Aquillaria agallocha* Roxh (*Agaru*), *Santalum alba* Linn. (*Hima*), *Coleus vettiveroides* KC Jacob. (*Ambu*), *Saussurea lappa* CB Cl (*Gada*), *Vetiveria zizinooides* Linn. (*Usheera*). The powder is usually mixed in hot water and applied over the joint region as a poultice.

Mechanical wear and tear is no longer considered the only cause of OA. Inflammation is an integral part of the disease process as seen necessitating the use of non-steroidal anti-inflammatory drugs (NSAIDs) frequently to manage the symptoms.^[4-5] Inflammation promotes synovitis and also enhances cartilage and bone destruction. The innate and adaptive immune systems are activated and pro-inflammatory cytokines are released [Charles J Malesud, 2015]. Prolonged use of NSAIDs however can adversely affect the liver, kidney, gut, skin and cardiovascular system and hence, has to be prescribed with extreme caution.^[6] As the disease affects the ageing population, comorbidities like hypertension, diabetes mellitus, cardiac disease etc. also prevent optimal prescription of oral medication. Hence safer and effective topical applications need to be explored as alternatives. Though there are few studies that have explored the potential of Ayurveda interventions in OA, most of them are restricted to evaluating the effect of oral medication. Herbs like *Tribulus terrestris* Linn. (*Gokshura*), *Tinospora cordifolia* Hook F & Thoms. (*Guduchi*), *Zingiber officinale* Rosc. (*Shunti*), *Emblia officinalis* Gaertn. (*Amalaki*), *Boswellia serrata* Roxh. (*Shallaki*), *Withania somnifera* Linn. (*Ashwagandha*) have been demonstrated to be safe and effective in ameliorating OA symptoms. These herbs have shown activity equivalent to that of standard care drugs like glucosamine sulphate.^[8-11]

A cursory review of topical herbal therapies for OA details Arnica gel, capsicum extract gel, comfrey extract gel and an adhesive patch of Chinese herbal mixture.^[13] An in vitro study of oral and topical application of boswellic acid showed reduction in cartilage loss, synovitis and osteophyte formation.^[14] Though topical applications are greatly used in Ayurveda for the management of OA, minimal data is currently available to demonstrate its effectiveness and also indicating the molecular and cellular mechanisms of their action. With this background, the anti-inflammatory effect of *Upanaha Choornam* (UC) was tested in mice induced with inflammation along with evaluating its activity on the expression of inflammatory markers.

MATERIALS AND METHODS

The polyherbal formulation UC was sourced from Vaidyaratnam Oushadhasala (P) Ltd., Thrissur, India, Batch no. 3843, manufactured in March 2015. The in-vitro and in-vivo study was conducted at Radiant Research Services Pvt. Ltd, Bangalore, India, following their Standard Operating Procedure.

In-vitro studies

The test systems used for in vitro evaluation were RAW 264.7 (macrophages of mouse) and L929 (connective tissue of mouse). The test product UC was stored under normal room temperature, coded as RR2706 in the records.

Preparation of Test Doses

For studies, each weighed UC were separately dissolved in distilled Dimethyl sulfoxide (DMSO) and volume was made up with Dulbecco's Modified Eagle Medium (DMEM) supplemented with 2% inactivated Fetal Bovine solution (FBS) to obtain a stock solution of 1 mg/ml concentration and sterilized by filtration. Serial two fold dilutions were prepared from this for carrying out cytotoxic studies.

Determination of Cytotoxicity in Raw and L929 cell lines^[15]

The monolayer cell culture was trypsinized and the cell count was adjusted to 1.0 x 10⁵ cells/ml using

DMEM containing 10% FBS. To each well of the 96 well microtitre plate, 0.1 ml of the diluted cell suspension (approximately 10,000 cells) was added. After 24 h, when a partial monolayer was formed, the supernatant was flicked off, washed the monolayer once with medium and 100 µl of different test concentrations of UC was added on to the partial monolayer in microtitre plates. The plates were then incubated at 37° C in 5% CO₂ atmosphere. After 24 h, the drug solutions in the wells were discarded and 50 µl of MTT in PBS was added to each well. The plates were gently shaken and incubated for 3 h at 37° C in 5% CO₂ atmosphere. The supernatant was removed and 100 µl of propanol was added and the plates were gently shaken to solubilize the formed formazan. The absorbance was measured using a microplate reader at a wavelength of 540 nm. The percentage growth inhibition was calculated and concentration of UC needed to inhibit cell growth by 50% (CTC₅₀) values is generated from the dose-response curves for each cell line.

In vitro TNF-α inhibitory activity of extracts^[16]

Step I: Induction of TNF-α in RAW cells: RAW cells seeded in to 6 well culture dishes at a cell population 1.5 to 2x10⁵ cells/ml in DMEM with 10% FBS. After 24 h, the cells were treated with known non-toxic concentration of test extracts along with 1µg/ml of lipopolysaccharide (LPS) and incubated at 37°C with 5% CO₂ for 4 h. After incubation, the cell supernatant was collected, centrifuged, separated and stored at -20° C till use.

Step II: Estimation of TNF-α in cell supernatant by bioassay: L929 monolayer cell culture was trypsinized and the cell count was adjusted to 1.0 x 10⁵ cells/ml using MEM medium containing 10% FBS. To each well of the 96 well microtitre plate, 0.1 ml of the diluted cell suspension (approximately 10,000 cells) was added. After 24 h, when a partial monolayer was formed, the supernatant was flicked off, washed the monolayer once with medium and 100 µl of diluted samples from the step I were added to the cells in quadruplicate wells. The cultures were then incubated at 37°C for 24 h in 5% CO₂ atmosphere. After 24h, the

drug solutions in the wells were discarded and 50 µl of MTT in PBS was added to each well. The plates were gently shaken and incubated for 3 h at 37°C in 5% CO₂ atmosphere. The supernatant was removed and 100 µl of propanol was added and the plates were gently shaken to solubilise the formed formazan. The absorbance was measured using a microplate reader at a wavelength of 540 nm and the percentage cell viability was calculated. The cell viability is direct indication of inhibitory properties of extracts against LPS induced TNF production in RAW cells.

Nitrite determination (Nitric oxide inhibition): RAW cells were treated with LPS and test samples as described above and incubated for 24 h and conditioned media collected were used for nitrite determination. Determination of nitrite as a biomarker for NO was carried out. In brief, equal volume (50 µL) of 0.1% N-1-naphthylethylenediamine dihydrochloride prepared in water, 1% sulfanilamide prepared in 5% phosphoric acid and cell culture media were mixed in flat bottom 96-well plate incubated for 10–15 min. Colored end product was measured at 530 nm. Percentage of nitric oxide inhibition was calculated over LPS control

RT-PCR Procedure: The mRNA expression levels of COX 2 carried out using semi-quantitative reverse transcriptase-polymerase chain reaction (RT-PCR). Briefly, the RAW cells were cultured in 60 mm petridish and maintained in DMEM medium for 48 hrs. The DMEM medium was supplemented with FBS and Amphotericin. To the dish was added the required concentration of Test sample (1000 and 500 µg/ml) and incubated for 24 hr. Total cellular RNA was isolated from the untreated (control) and treated cells using Tri Reagent according to manufacturer's protocol. cDNA was synthesized from total isolated RNA by reverse transcriptase kit according to manufacturer's instructions. Then 50µl of the reaction mixture was subjected to PCR for amplification of COX 2 cDNAs using specifically designed primers procured from Eurofins India, as an internal control, the house keeping gene GAPDH was co-amplified with each reaction.

In vivo study

The in vivo study was performed in Swiss Albino mice (n=24) of either gender, by TPA induced ear oedema. The study was approved by committee for the purpose of control and supervision of experiments on animals (CPCSEA) and Institutional Animal Ethics Committee (IAEC) with registration number 1803/PO/RcBiS/2015/CPCSEA.

The mice were randomly grouped into 4 groups with 6 mice in each, weighing 25 to 30 g. accommodated in standard polycarbonate cages under standard laboratory conditions. The animals were acclimatised using laboratory conditions for 5 days, 7 days and 9 days respectively for step I, step II and step III respectively after veterinary examination. Only the animals without any physical signs of illness were selected. The mode of application of the UC was topical. The animals were allocated as: Group I was Control group received normal saline only, Group II was treatment group treated with STD (Indomethacin 0.5mg), Group III was Treatment group treated with UC (30% of UC) and Group IV was treatment group treated with UC (60% of UC). Thirty percent is 1ml Tween 80 + 300mg UC, sixty percent is 1ml Tween 80 + 600mg UC (Tween 80 was procured from Central Drug House, Bangalore).

The left ear was considered as contra lateral control and received only saline solution. Inflammation was induced to the ear of mice by applying TPA (12-O-tetradecanoyl-phorbol-13-acetate, procured from Sigma, Bangalore) (2.5 µg/ear) and Xylene (20 µL/ear) on the inner and outer surfaces of the right ear and left undisturbed for 30 mins. Application was done between 9.00 and 9.30 am and ear oedema was evaluated at 0, 30, 60, 120, 180, 240 and 300 mins after TPA application. After 5 hours of the treatment, mice were subjected to CO₂ euthanasia and sacrificed by cervical dislocation and each ear (8mm diameter) biopsies was done and weighed immediately. Oedema was measured in terms of weight difference of the ears and the ear thickness was measured before and after induction of the inflammatory response using a digital micrometer.

STATISTICAL ANALYSIS

The In vivo data are represented as Mean ± SE. Comparison between groups was made by One-way ANOVA, differences with p<0.05 were considered significant.

RESULTS

In vitro study: UC showed moderate cytotoxicity, on the basis of CTC₅₀ value the concentration of the UC in RAW cell line. The cytotoxicity activity against the RAW cells and L929 cells increased with increasing dose (Tables 1-2). The test product showed dose dependent anti-inflammatory activity with moderate TNF-alpha inhibition, nitric oxide inhibition properties (Tables 3-4). In gene expression study, UC showed regulation on the levels of COX-2 expression. UC i.e. 1000 µg/ml and 500 µg/ml up regulate the COX-2 level by 0.08 fold and 0.03 as compare to control (Figure 1).

In vivo study: UC decreased the TPA induced ear oedema during the observation period when compared to control group (F=1250, p<0.001) (Figure 2). The inhibition was found to be 52.3 and 61.9 % at 300 mins, respectively (Table 5).

Table 1: Cytotoxic properties of UC against RAW cell line

Test Conc. (%)	% Cytotoxicity	CTC ₅₀ (µg/ml)
1000	44.91±1.2	>1000
500	43.59±1.8	
250	30.73±0.5	
125	27.22±2.9	
62.5	15.24±1.3	

Table 2: Cytotoxic properties of UC against L929 cell line

Test Conc. (%)	% Cytotoxicity	CTC ₅₀ (µg/ml)
1000	43.83±1.1	>1000

500	36.62±2.5	
250	26.19±1.2	
125	16.71±1.8	
62.5	7.05±1.2	

Table 3: Anti-inflammatory effect of UC in RAW cells against LPS induced TNF production

SN	Material	Concentration tested	% TNF inhibition
1	UC	500 µg/ml	48.2±1.9
2	UC	250 µg/ml	25.6±1.0
3	Dexamethasone	200 µM	67.3±1.7

Table 4: Anti-inflammatory effect of UC in RAW cells against LPS induced Nitric oxide production.

SN	Concentration tested	% NO inhibition
1	500 µg/ml	14.8±0.39
2	250 µg/ml	9.5±0.13

Table 5: Effect of UC on TPA induced ear edema in mice

GRO UPS	Ear thickness (mm)						
	0 mins	30 mins	60 mins	120 mins	240 mins	300 mins	% Inhibition
Group I	0.19	0.31	0.37	0.40	0.41	0.40	-
Group II	0.24	0.34	0.32	0.29	0.27	0.26	90.4
Group III	0.23	0.37	0.37	0.35	0.34	0.33	52.3
Group IV	0.21	0.32	0.33	0.31	0.30	0.29	61.9

Figure 1




Figure 2





Figure 3



DISCUSSION

Targeting inflammation is considered as the initial step for effective management of Osteoarthritis. In the current study, the anti-inflammatory potential of

an Ayurveda polyherbal formulation *Upanaha Churna* (UC) used widely for topical application in *Sandhigata Vata* (OA) was evaluated. Both in vitro and in vivo experimental protocols were followed and significant effects were obtained. The study has aided understanding of the probable mode of action of the polyherbal formulation UC.

To the best of our knowledge, there are very few studies that have looked into the utilitarian aspect of topical applications for inflammatory conditions like OA. The latest recommendations for OA management too indicate towards the positive outcomes of topical applications.^[18] The mechanism of action by which these provide analgesia is proposed to be similar to oral medications, albeit with a better safety profile.^[1] Topical NSAIDs have produced fewer systemic adverse events and also shown effect in OA as per a Cochrane review.^[19] Though even in Ayurveda pharmacopoeia, topical applications are widely used for *Sandhigata Vata* (OA), there are very few studies that have delved into the mechanisms by which these medications work. Singh and colleagues reported the anti-inflammatory activity of Boswellic acids when applied topically in models of acute and chronic inflammation.^[20] Another in vitro study of a polyherbal formulation containing *Z. officinalis*, *T. cordifolia*, *E.officinalis* and *B.serrata* on human knee cartilage has shown significant reduction of breakdown products glycosaminoglycan and aggrecan.^[21] The chondroprotective effect and regenerative ability of *Cissus quadrangularis* Linn. (*Asthishrinkala*) has also been demonstrated in human chondrocytes. The effect is attributed to the inhibition of the p38MAPK signalling pathway by the upregulation of *survivin*.^[22] A polyherbal compound BV-9238 containing *W. somnifera*, *B. serrata*, *Z. officinale* and *C. longa* inhibited TNF- α and nitric oxide (NO) production in RAW cell lines.^[23] Curcumin, by virtue of its anti-apoptotic and anti-catabolic effects on IL-1 β -stimulated articular chondrocytes, is also considered as a potent remedy in OA.^[24] It also inhibits COX-2 activity.^[25]

In clinical settings too, herbal topical applications have shown reduction in pain and stiffness, and

improvement in mobility. A cream containing *Trigonella foenum-graceum* Linn. (fenugreek), *Curcuma longa* Linn. (turmeric), coconut oil, castor oil, capsaicin, *Mentha sylvestris* Linn. (menthol) and other ingredients reduced pain and stiffness in hand and knee OA. The activity of the cream though decayed at onset, sustained for longer duration when compared with other topical applications.^[26] A meta-analysis of topical herbal applications reported that trials of boswellia and curcumin mixture and *B.serrata* alone, reduced pain and improved movement.^[27] Another herbal ointment containing *Cinnamomum verum* Presl. (cinnamon), *G. officinalis*, *saghez* (resin of tree *Pistacia lentiscus*) and sesame oil decreased pain, morning stiffness and limited motion in patients of knee OA; its effect were comparable with Salicylate ointment.^[28] Topical ginger (*Z. officinalis*) compress and pack reduced pain and fatigue and improved functionality and satisfaction in OA patients.^[29] Herbal compounds are hypothesized to modulate several inflammation triggering factors, pathway mediators and conditions, thereby offering relief in such conditions.^[30]

In the current study, *Upanaha Choornam* showed a significant anti-inflammatory activity, comparable to that of standard analgesic Indomethacin. Increased concentration did not seem to offer better benefit. The in-vitro studies helped to understand the mechanism of action of the polyherbal formulation. The role of inflammatory cytokines in the pathogenesis of OA is seemingly very clear. TNF- α and IL-1 β are prominently involved, the former driving the inflammatory process and the latter contributing to cartilage destruction.^[31] Cartilage breakdown is also promoted by NO, though its beneficial effects are also proposed.^[32] Increase in COX-2 levels is also observed in inflammation, leading to efforts to curb its expression.^[33] It is also proposed that anti-cytokine therapy may be a potential Disease-modifying OA drugs.^[34-36] UC too inhibited TNF- α and NO and COX-2 expression and hence we may infer that it has the potential to reduce destruction in OA.

Herbs have been used since times immemorial for their medicinal properties, especially in India which

has its own indigenous healthcare system Ayurveda. We have here reported the anti-inflammatory activity of an Ayurveda polyherbal formulation *Upanaha Chooranam*, used as a topical application in Osteoarthritis. This may have implications in better management of this degenerative condition.

REFERENCES

- Balmaceda CM. Evolving guidelines in the use of topical nonsteroidal anti-inflammatory drugs in the treatment of osteoarthritis. *BMC Musculoskeletal Disorders* 2014; 15:27.
- Cross M, Smith E, Hoy D, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014; 73:1323-30.
- Wittenauer R, Smith L, Aden K. Background Paper 6.12 Osteoarthritis. World Health Organization.
- Available from: http://www.who.int/medicines/areas/priority_medicines/BP6_12Osteo.pdf.
- Hochberg MC. Osteoarthritis year 2012 in review: clinical. *Osteoarthritis Cartilage* 2012; 20:1465-9.
- Zhang W, Moskowitz RW, Nuki G, et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage* 2008; 16:137-62.
- Scarpignato C, Lanus A, Blandizzi C, et al. Safe prescribing of non-steroidal anti-inflammatory drugs in patients with osteoarthritis--an expert consensus addressing benefits as well as gastrointestinal and cardiovascular risks. *BMC Med* 2015; 13:55.
- Charaka, Drdhabala, Chakrapanidatta. The Charaka samhita of Agnivesha. 5 ed. Varanasi: Munshiram Manoharlal Publishers Pvt. Ltd; 1992.
- Chopra A, Saluja M, Tillu G, et al. Ayurvedic medicine offers a good alternative to glucosamine and celecoxib in the treatment of symptomatic knee osteoarthritis: a randomized, double-blind, controlled equivalence drug trial. *Rheumatology (Oxford)* 2013; 52:1408-17.
- Chopra A, Saluja M, Tillu G, et al. A Randomized Controlled Exploratory Evaluation of Standardized Ayurvedic Formulations in Symptomatic Osteoarthritis Knees: A Government of India NMITLI Project. *Evid Based Complement Alternat Med* 2011; 2011:724291.
- Nipanikar SU, Saluja M, Kuber VV, Kadbhane KP, Chopra A, Khade NR. An open label, prospective, clinical study on a polyherbal formulation in osteoarthritis of knee. *J Ayurveda Integr Med* 2013; 4:33-9.
- Raut A, Bichile L, Chopra A, Patwardhan B, Vaidya A. Comparative study of amrutbhallataka and glucosamine sulphate in osteoarthritis: Six months open label randomized controlled clinical trial. *J Ayurveda Integr Med* 2013; 4:229-36.
- Vagbhata, Arunadatta, Hemadri. *Ashtanga hridaya*. Varanasi: Krishnadas Academy; 1995.
- Cameron M, Chrubasik S. Topical herbal therapies for treating osteoarthritis. *Cochrane Database System Review* 2013; 5:CD010538.
- Wang Q, Pan X, Wong HH, et al. Oral and topical boswellic acid attenuates mouse osteoarthritis. *Osteoarthritis Cartilage* 2014; 22:128-32.
- Denizot F, Lang R. Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *Journal of Immunological Methods* 1986; 89:271-7.
- Varma RS, Ashok G, Vidyashankar S, Patki P, Nandakumar KS. Anti-inflammatory properties of Septilin in lipopolysaccharide activated monocytes and macrophage. *Immunopharmacol Immunotoxicol* 2011; 33:55-63.
- Choudhary M, Kumar V, Malhotra H, Singh S. Medicinal plants with potential anti-arthritic activity. *Journal of Intercultural Ethnopharmacology* 2015; 4:147-79.
- Bruyere O, Cooper C, Pelletier JP, et al. An algorithm recommendation for the management of knee osteoarthritis in Europe and internationally: a report from a task force of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Semin Arthritis Rheum* 2014; 44:253-63.
- Derry S, Moore RA, Rabbie R. Topical NSAIDs for chronic musculoskeletal pain in adults. *Cochrane Database Syst Rev* 2012; 9:CD007400.
- Singh S, Khajuria A, Taneja SC, Johri RK, Singh J, Qazi GN. Boswellic acids: A leukotriene inhibitor also

- effective through topical application in inflammatory disorders. *Phytomedicine* 2008; 15:400-7.
22. Sumantran VN, Joshi AK, Boddul S, et al. Antiarthritic activity of a standardized, multiherbal, Ayurvedic formulation containing *Boswellia serrata*: in vitro studies on knee cartilage from osteoarthritis patients. *Phytother Res* 2011; 25:1375-80.
 23. Kanwar JR, Samarasinghe RM, Kumar K, et al. *Cissus quadrangularis* inhibits IL-1 β induced inflammatory responses on chondrocytes and alleviates bone deterioration in osteotomized rats via p38 MAPK signaling. *Drug Des Devel Ther* 2015; 9:2927-40.
 24. Dey D, Chaskar S, Athavale N, Chitre D. Inhibition of LPS-induced TNF- α and NO production in mouse macrophage and inflammatory response in rat animal models by a novel Ayurvedic formulation, BV-9238. *Phytotherapy Research*, 2014; 28:1479-85.
 25. Shakibaei M, Schulze-Tanzil G, John T, Mobasheri A. Curcumin protects human chondrocytes from IL-1 β -induced inhibition of collagen type II and β 1-integrin expression and activation of caspase-3: An immunomorphological study. *Ann Anat* 2005; 187:487-97.
 26. Lev-Ari S, Strier L, Kazanov D, et al. Curcumin synergistically potentiates the growth-inhibitory and pro-apoptotic effects of celecoxib in osteoarthritis synovial adherent cells. *Rheumatology (Oxford)* 2006; 45:171-7.
 27. Gemmell HA, Jacobson BH, Hayes BM. Effect of a topical herbal cream on osteoarthritis of the hand and knee: a pilot study. *J Manipulative Physiol Ther* 2003; 26:e15.
 28. Grover AK, Samson SE. Benefits of antioxidant supplements for knee osteoarthritis: rationale and reality. *Nutr J* 2016; 15:1.
 29. Zahmatkash M, Vafaenasab MR. Comparing analgesic effects of a topical herbal mixed medicine with salicylate in patients with knee osteoarthritis. *Pak J Biol Sci* 2011; 14:715-9.
 30. Therkleson T. Topical Ginger Treatment with a Compressor Patch for Osteoarthritis Symptoms. *J Holist Nurs* 2013; 32:173-82.
 31. Maroon JC, Bost JW, Maroon A. Natural anti-inflammatory agents for pain relief. *Surg Neurol Int* 2010; 1:80.
 32. Martel-Pelletier J, Alaaeddine N, Pelletier JP. Cytokines and their role in the pathophysiology of osteoarthritis. *Front Biosci* 1999; 4:D694-703.
 33. Abramson SB. Osteoarthritis and nitric oxide. *Osteoarthritis Cartilage* 2008; 16 Suppl 2:S15-20.
 34. Rao P, Knaus EE. Evolution of nonsteroidal anti-inflammatory drugs (NSAIDs): cyclooxygenase (COX) inhibition and beyond. *J Pharm Pharm Sci* 2008; 11:81s-110s.
 35. Kapoor M, Martel-Pelletier J, Lajeunesse D, Pelletier JP, Fahmi H. Role of proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat Rev Rheumatol* 2011; 7:33-42.
 36. Malemud CJ. Anticytokine therapy for osteoarthritis: evidence to date. *Drugs Aging* 2010; 27:95-115.
 37. Wojdasiewicz P, Poniatowski LA, Szukiewicz D. The role of inflammatory and anti-inflammatory cytokines in the pathogenesis of osteoarthritis. *Mediators Inflamm* 2014; 2014:561459.

How to cite this article: Sitaram Ahalya, B. A. Venkatesh, R. Vijayasarithi, Tirumalapura Vijayanna Shalini. Anti-inflammatory effect of an Ayurveda polyherbal topical application and exploration of its mechanism of action. *J Ayurveda Integr Med Sci* 2017;6:34-41.

<http://dx.doi.org/10.21760/jaims.v2i06.10924>

Source of Support: Nil, **Conflict of Interest:** None declared.
